

# PILOT AWARDS

Developing and submitting a  
competitive, successful application

---

Tammy Hshieh, MD MPH



NIDUS

Network for Investigation of  
Delirium: Unifying Scientists

# My Project

- Eventually:
  - Delirium severity and long-term clinical outcomes among persons with and without Alzheimer's Disease and Related Dementias
- For the pilot:
  - Feasibility of assessing delirium severity
  - Feasibility of determining who has ADRD

# Scope

- What is in scope?
  - Deciding which delirium, delirium severity instruments to use
  - Deciding which measure of delirium severity to use in analysis
  - Defining ADRD for this study population
- What is out of scope?
  - Clinical outcomes (function, nursing home placement, length of stay, re-hospitalization)

# Refining methods

- Cognitive status → Brief cognitive testing
- Delirium → Confusion Assessment Method (CAM)
- Delirium severity → CAM-Severity (CAM-S)
  - Peak CAM-S
  - Sum of all CAM-S scores
  - Categorical vs. continuous

# Refining methods

- ADRD status determined by cognitive testing, chart review and expert consensus
  - Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE)  $\geq 3.2$
  - Documentation of dementia by ICD-10 codes
  - Impairment on any of three cognitive tests: Montreal Cognitive Assessment  $< 26$ , brief cognitive screen  $< 14/16$  or MoCA delayed recall of  $\leq 2$
  - Very variable documentation of cognitive impairment

# Refining methods

- Of the 352 original patients, 197 participants needed chart review by expert clinicians
  - Medications (cholinesterase inhibitors, memantine, vitamin E)
  - Neuroimaging studies
  - Provider notes (inpatient and outpatient) +/- formal neuropsychological evaluation
  - For 24 cases with limited data or lack of agreement, adjudicated by interdisciplinary expert panel until consensus was reached

# Outcomes

- Used peak CAM-S categorically
  - Delirium incidence:
    - Persons with ADRD = 45%
    - Persons without ADRD = 19% (50/267)
- ADRD patients
  - 24% of cohort (85/352)
  - Higher peak CAM-S scores

# Next steps

- We need a tool that better assesses delirium severity in ADRD patients!
  - External and local expert panels
  - DEL-S-AD
- We need to incorporate more ADRD patients into clinical studies



# Thank you!

- Questions?
- [thshieh@bwh.harvard.edu](mailto:thshieh@bwh.harvard.edu)

